

Samples	MPA 61285	MPA 61286	MPA 61287	MPA 61288
Fractions (µm)	500	500	500	500
<i>Acroloxus lacustris</i> (Linné, 1758) % of fraction	5 10.63			1 4.54
<i>Bathynomphalus contortus</i> (Linné, 1758) % of fraction				1 4.54
<i>Bithynia tentaculata</i> (Linné, 1758) % of fraction		5 5.81		1 4.54
<i>Bithynia leachii</i> (Sheppard, 1823) % of fraction		6 6.97		
<i>Pisidium amnicum</i> (Müller, 1774) % of fraction		1 1.16		
<i>Pisidium henslowanum</i> (Sheppard, 1823) % of fraction	4 8.51	6 6.97		
<i>P. Moitessierianum</i> Paladilhe, 1866 % of fraction	3 6.38		4 40	1 4.54
<i>P. nitidum</i> Jenyns, 1832 % of fraction	13 27.65	11 12.79	1 10	3 13.63
? <i>P. pseudospaerium</i> ? (Favre, 1927) % of fraction	2 4.25			
<i>Planorbis carinatus</i> Müller, 1774 % of fraction	1 2.12	7 8.13		
<i>Radix balthica</i> (Müller, 1774) % of fraction		3 3.48		
<i>Radix labiata</i> (Rossmässler, 1835) % of fraction	5 10.63		1 10	
<i>Unio tumidus</i> Philipsson, 1788 % of fraction			1 10	
<i>Valvata cristata</i> (Müller, 1774) % of fraction	1 23.40	14 16.27		3 13.63
<i>Valvata macrostoma</i> Mörch, 1864 % of fraction	1 2.12	20 23.25	2 20	5 22.72
<i>V. piscinalis</i> (Müller, 1774) % of fraction	2 4.25	2 2.32		4 18.18
<i>Viviparus viviparus</i> (Linné, 1758) % of fraction		11 12.79	1 10	3 13.63
Total %total	47 100	86 100	10 100	22 100